

RESEARCH

Open Access

# A mixture of anatase and rutile TiO<sub>2</sub> nanoparticles induces histamine secretion in mast cells

Eric Y Chen<sup>†</sup>, Maria Garnica<sup>†</sup>, Yung-Chen Wang, Alexander J Mintz, Chi-Shuo Chen and Wei-Chun Chin\*

## Abstract

**Background:** Histamine released from mast cells, through complex interactions involving the binding of IgE to FcεRI receptors and the subsequent intracellular Ca<sup>2+</sup> signaling, can mediate many allergic/inflammatory responses. The possibility of titanium dioxide nanoparticles (TiO<sub>2</sub> NPs), a nanomaterial pervasively used in nanotechnology and pharmaceutical industries, to directly induce histamine secretion without prior allergen sensitization has remained uncertain.

**Results:** TiO<sub>2</sub> NP exposure increased both histamine secretion and cytosolic Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>C</sub>) in a dose dependent manner in rat RBL-2H3 mast cells. The increase in intracellular Ca<sup>2+</sup> levels resulted primarily from an extracellular Ca<sup>2+</sup> influx via membrane L-type Ca<sup>2+</sup> channels. Unspecific Ca<sup>2+</sup> entry via TiO<sub>2</sub> NP-instigated membrane disruption was demonstrated with the intracellular leakage of a fluorescent calcein dye. Oxidative stress induced by TiO<sub>2</sub> NPs also contributed to cytosolic Ca<sup>2+</sup> signaling. The PLC-IP<sub>3</sub>-IP<sub>3</sub> receptor pathways and endoplasmic reticulum (ER) were responsible for the sustained elevation of [Ca<sup>2+</sup>]<sub>C</sub> and histamine secretion.

**Conclusion:** Our data suggests that systemic circulation of NPs may prompt histamine release at different locales causing abnormal inflammatory diseases. This study provides a novel mechanistic link between environmental TiO<sub>2</sub> NP exposure and allergen-independent histamine release that can exacerbate manifestations of multiple allergic responses.

**Keywords:** TiO<sub>2</sub> nanoparticles, mast cell, histamine release, Ca<sup>2+</sup> signaling

## Background

Allergic inflammation is a primary pathological feature of many debilitating diseases [1]. Among the numerous active mediators and cytokines that modulate initiation and progression of allergic inflammation, histamine is distinctly potent [1,2]. Typically, the storage of histamine is restricted to mast cells and circulating basophils [2,3]. The cardinal pathway of histamine release involves the attachment of IgE-bound allergens to high-affinity FcεRI receptors on mast cells and the crosslinking of adjacent IgE molecules by allergens [1,2]. Subsequent receptor clustering leads to a complex cascade of intracellular Ca<sup>2+</sup> signaling resulting from increased activity of phospholipase C (PLC), generation of diacylglycerol (DAG) (activating PKC) and inositol 1,4,5-trisphosphate (IP<sub>3</sub>) which mobilizes the ER Ca<sup>2+</sup> store and participates in final histamine

secretion from mast cells. Activation of histamine receptors (H1, H2, H3 and H4) greatly influences inflammatory responses [2].

Aside from inducing acute allergic inflammatory responses, histamine also mediates chronic phase progression by augmenting the secretion of pro-inflammatory cytokines such as IL-1α, IL-1β, and IL-6 as well as chemokines like RANTES [1,4]. Consequential pathologies are expressed in many systems encompassing ocular, airway, skin and GI tracts [2]. Associated disorders may include asthma, allergic rhinitis, allergic conjunctivitis, atopic dermatitis, urticaria, anaphylaxis and food allergies [1,4,5]. Possible clinical symptoms include itchiness, increased vascular permeability, edema, leukocyte infiltration, bronchoconstriction and mucus hypersecretion [1,4,5]. Therefore, any disturbance to the immunological and/or homeostatic control of histamine release can potentially intensify inflammation leading to health problems.

\* Correspondence: wchin2@ucmerced.edu

† Contributed equally

Bioengineering, University of California at Merced, Merced, CA, USA. 5200 North Lake RD, Merced, CA 95343, USA

Recently, numerous epidemiological studies have suggested that pollution associated airborne particulate matter (PM) can aggravate allergic inflammatory responses. Classical examples of allergies indicate that people with asthma and rhinitis are more susceptible to the short term acute effects of particle exposure [6-8]. In line with epidemiological studies, results from animal models also demonstrated that ultrafine particles (one of the major components of PM) can modulate asthmatic responses by exacerbating pulmonary inflammation and airway hyper-responsiveness [9-11]. In an atopic, dermatitis-like, skin lesion mouse model, exposure to titanium dioxide nanoparticles (TiO<sub>2</sub> NPs) was found to worsen symptoms by elevating pro-inflammatory molecules in the skin and increasing serum levels of IgE and histamine [12]. At the same time, exposure to environmental tobacco smoke has been found to increase the risks of rhinoconjunctivitis and allergic conjunctivitis [13,14]. Moreover, Kulig et al reported that pre-natal and postnatal exposure to environmental tobacco smoke in children (< 3 yrs old) was associated with sensitization to food allergens [15]. Despite NPs' ability to potentiate allergic responses, the role that histamine plays in mediation is not clear. More importantly, whether NPs can directly modulate histamine release from mast cells without allergen sensitization remains elusive.

TiO<sub>2</sub> NPs have been extensively utilized in the nanotechnological and pharmaceutical arenas and are one of the main components in many household commodities and personalized products [16,17]. The enormous annual global production of TiO<sub>2</sub> broadens the possibilities of occupational and environmental exposures [18]. As a common constituent of PM<sub>10</sub>, TiO<sub>2</sub> NPs are widely known for their potential hazardous effects, which manifest biologically via inflammatory responses [19]. TiO<sub>2</sub> NPs size, surface area and crystalline structure ascribe cellular nanotoxicity [20-23]. In addition to respiratory inhalation, NPs can enter the human body via alternative routes such as: direct penetration through skin, ingestion and injection [24]. Animal studies have shown that intra-tracheally instilled TiO<sub>2</sub> and other NPs can possibly transmigrate from lung to systemic circulation [25-27]. Upon entering blood circulation, NPs can infiltrate multiple organs, potentially directly stimulating mast cells, a critical effector, exacerbating pathological consequences [24]. As such, TiO<sub>2</sub> NPs were selected as ideal model particles for our study.

Previously, we have demonstrated that TiO<sub>2</sub> NPs can trigger a cascade of cytosolic Ca<sup>2+</sup> signaling leading to mucin secretion [17]. In the present study, we aim to investigate the impact of NPs on histamine secretion from RBL-2H3 mast cells. We hypothesize that TiO<sub>2</sub> NPs can directly induce histamine release without prior allergen sensitization via a Ca<sup>2+</sup>-mediated pathway.

## Results

### TiO<sub>2</sub> NP characterization


Dynamic laser scattering (DLS) was used to characterize the TiO<sub>2</sub> NPs in suspension. The particle size in Hanks' solution had a distribution of ~6 to 100 nm due to minor aggregation while the predominant size was ~83 nm (Figure 1A). Transmission electron microscopy (TEM) provided detailed characterization of TiO<sub>2</sub> NP size where the mean particle diameter was found to be 60 ± 10 nm (Figures 1B and 1C).

### TiO<sub>2</sub> NP-Induced [Ca<sup>2+</sup>]<sub>C</sub> increase

We tested whether TiO<sub>2</sub> NPs could trigger [Ca<sup>2+</sup>]<sub>C</sub> increase by loading RBL-2H3 cells with Rhod-2 AM dye and exposing them to 0.1 mg/ml-1 mg/ml of TiO<sub>2</sub> NPs. The TiO<sub>2</sub> concentration range used in our study is consistent with the concentrations found in ambient, nanotechnology industries and existing literatures [17,20,28]. The changes in [Ca<sup>2+</sup>]<sub>C</sub> were measured by monitoring the intracellular fluorescence intensity. Figure 2A shows that 1 mg/ml induced an approximate 160% increase, while lower TiO<sub>2</sub> NP concentrations (< 0.25 mg/ml) caused a smaller elevation, when compared to untreated cells. The results demonstrated a TiO<sub>2</sub> NP concentration dependent increase in [Ca<sup>2+</sup>]<sub>C</sub> (Figure 2A).

### Extracellular source of [Ca<sup>2+</sup>]<sub>C</sub> increase

The source of elevated [Ca<sup>2+</sup>]<sub>C</sub> was identified by stimulating RBL-2H3 cells with TiO<sub>2</sub> NPs in Ca<sup>2+</sup>-free Hanks' buffer with EGTA added to chelate traces of Ca<sup>2+</sup>. Figure 2B demonstrates that TiO<sub>2</sub> NPs (0.1 mg/ml-1 mg/ml) failed to induce a significant increase in [Ca<sup>2+</sup>]<sub>C</sub>, when compared with the 160% increase observed in normal Hanks' buffer (Figure 2A). Our data suggested that the extracellular Ca<sup>2+</sup> pool was the primary source of the observed [Ca<sup>2+</sup>]<sub>C</sub> increase. We then examined if TiO<sub>2</sub> NPs can induce extracellular Ca<sup>2+</sup> influx via membrane channels, in particular L-type Ca<sup>2+</sup> channels. Blocking the channels with CdCl<sub>2</sub> (200 μM) markedly inhibited the rise in [Ca<sup>2+</sup>]<sub>C</sub> by approximately 75% (Figure 2C). Subsequent pre-treatment of cells with nifedipine or verapamil (L-type Ca<sup>2+</sup> channel blockers) also hampered the rise in [Ca<sup>2+</sup>]<sub>C</sub> elicited by TiO<sub>2</sub> NPs (Figure 2D and 2E). Nonetheless, the incomplete blockage of Ca<sup>2+</sup> influx with membrane channel blockers suggests a possible Ca<sup>2+</sup> leakage through perturbed cell membranes. To confirm that TiO<sub>2</sub> NPs can instigate membrane disruption, thereby permitting unspecific extracellular Ca<sup>2+</sup> entry, a fluorescent calcein dye was used to assess cytosolic leakage. Results showed a dye permeation ratio increase from approximately 5 to 12% with a TiO<sub>2</sub> NP concentration ranging from 0.1 to 1 mg/ml (Figure 2F).



### Induction of oxidative stress and associated Ca<sup>2+</sup> influx

Demonstrating possible effects of oxidative stress evoked by TiO<sub>2</sub> NPs, intracellular ROS formation was investigated. Transient exposure of RBL-2H3 cells to TiO<sub>2</sub> NPs for 15 min resulted in cytosolic ROS level increase in a dose-dependent manner (Figure 3A). Since oxidative stress can trigger intracellular Ca<sup>2+</sup> signaling, cells pretreated with an antioxidant NAC were monitored for changes in [Ca<sup>2+</sup>]<sub>C</sub> upon TiO<sub>2</sub> stimulation. Our data revealed that NAC notably attenuated the increase in [Ca<sup>2+</sup>]<sub>C</sub> triggered by TiO<sub>2</sub> NPs (Figure 3B).

### The IP<sub>3</sub>-IP<sub>3</sub> receptor as a histamine-release pathway

To understand the influence of PLC, and the ensuing IP<sub>3</sub>-IP<sub>3</sub> receptor pathway on cytosolic Ca<sup>2+</sup>, the RBL-2H3 cells were pre-treated with U73122 (PLC inhibitor), 2-APB or Xestospongine (IP<sub>3</sub> receptor blockers), before being exposed to TiO<sub>2</sub> NPs. The data revealed a > 75% drop in the [Ca<sup>2+</sup>]<sub>C</sub> after TiO<sub>2</sub> NP stimulation (Figure 4A-C). Since ER is one of the main intracellular Ca<sup>2+</sup> stores, it is highly responsible for the amplification of intracellular Ca<sup>2+</sup> signals. Thapsigargin was used to deplete intracellular ER Ca<sup>2+</sup> pool and caused a decrease in the [Ca<sup>2+</sup>]<sub>C</sub> after TiO<sub>2</sub> NP exposure (Figure 4D).


### Ca<sup>2+</sup> dependency of TiO<sub>2</sub> NP-induced histamine secretion

ELISA was used to assess the amount of histamine secreted from RBL-2H3 cells when stimulated with TiO<sub>2</sub> NPs. Comparing with the control, TiO<sub>2</sub> NPs increased histamine secretion in a dose-dependent fashion (Figure 5A). Incubation with BAPTA-AM (intracellular calcium chelator), Thapsigargin, U73122, 2-APB or Xestospongine significantly attenuated histamine secretion (Figure 5B). Our data indicated that histamine release is attributed to a high [Ca<sup>2+</sup>]<sub>C</sub> sustained by an external Ca<sup>2+</sup> influx and a secondary Ca<sup>2+</sup> release from intracellular Ca<sup>2+</sup> stores (organelles).


### Discussion

Recently, a growing epidemic of exacerbated allergic inflammatory diseases have been closely linked with the exposure to airborne PM pollution [5,7,8]. Many prior investigations have focused on the hazardous effects of NPs. These reports have primarily been either epidemiological studies that draw correlations between clinical manifestations and exposure to environmental pollutants or immunological studies that examine the release of inflammatory cytokines and subsequent recruitment of immune cells using allergen-sensitized mice with PM or NP insults [7,9,10]. Few studies have systematically investigated the role of histamine secretion from mast cells during NP challenge and its role in mediating allergic inflammation. By using allergen-sensitized animal models, these studies overlooked the NP's potential to directly stimulate histamine release from mast cells without prior allergen sensitization or IgE production. In addition, the associated underlying cellular mechanisms leading to the final degranulation are not clear. TiO<sub>2</sub> NPs have previously been shown to trigger mucin secretion, pulmonary inflammatory responses and emphysema-like pathology [17,18,29,30]. In this study, we demonstrate that TiO<sub>2</sub> NPs can directly stimulate histamine release from RBL-2H3 mast cells via a Ca<sup>2+</sup>-dependent pathway.

Allergic inflammatory diseases encompass a multifactorial interplay between many elements/systems; our study focuses on how TiO<sub>2</sub> NPs can directly trigger histamine release from mast cells and its potential to exacerbate allergic symptoms. This effect is due to histamine's ability to affect the physiology and pathology of a wide range of cells via corresponding receptors that are widely expressed on airway and vascular muscle cells, hepatocytes, chondrocytes, endothelial cells, epithelial cells, neutrophils, eosinophils, monocytes, macrophages, dendritic cells, and T and B cells [4]. We found that TiO<sub>2</sub> NPs can significantly



**Figure 2 Measurement of the  $[\text{Ca}^{2+}]_c$  and calcein leakage after  $\text{TiO}_2$  NP exposure.** (A) RBL-2H3 cells were stimulated with  $\text{TiO}_2$  NPs with concentrations of 0.1 mg/ml (orange), 0.25 mg/ml (green), 0.5 mg/ml (purple), 0.75 mg/ml (blue) and 1 mg/ml (red) in normal Hanks' solution, (B) in  $\text{Ca}^{2+}$ -free Hanks' solution, (C) in the presence of  $\text{CdCl}_2$  (200  $\mu\text{M}$ ), (D) nifedipine (10  $\mu\text{M}$ ), (E) verapamil (100  $\mu\text{M}$ ), and (F) calcein (50  $\mu\text{M}$ ) ( $n \geq 12$ ,  $**p < 0.005$ ) (all colors are as depicted in Figure 2A and each line is a representative fluorescence intensity of approximately 200 cells).




**Figure 3** Measurement of intracellular ROS level and the associated changes in  $[Ca^{2+}]_c$ . (A) RBL-2H3 cells were exposed to  $TiO_2$  NPs at concentrations 0, 0.1, 0.25, 0.5 and 0.75 mg/ml. The generation of ROS was measured by fluorescence imaging. The level of ROS increased as a function of increasing  $TiO_2$  NP concentration ( $n \geq 50$ ,  $**p < 0.005$ ). (B) Pretreatment with antioxidant NAC (250  $\mu M$ ) significantly reduced the rise in cytosolic  $Ca^{2+}$  when stimulated with  $TiO_2$  NPs. Each line is a representative fluorescence intensity of approximately 200 cells and the colors used are consistent with Figure 2A.

induce histamine release from RBL-2H3 cells in a dose dependent manner (Figure 5A). Our data is corroborated by evidence documenting more severe  $TiO_2$  NP-induced inflammation with elevated release of inflammatory cytokines than that of  $TiO_2$  of micrometric dimensions [31,32]. The  $TiO_2$  NP-elicited mast cell degranulation also appeared to be a  $Ca^{2+}$ -dependent process, as indicated by a major attenuation in histamine secretion when pretreated with BAPTA (Figure 5B). The importance of  $Ca^{2+}$  in stimulus-secretion coupling mechanisms is well documented [33,34]. We and others have reported that NPs increase  $[Ca^{2+}]_c$  prior to cellular exocytosis [17,35]. As  $TiO_2$  concentrations increased from 0.1 to 1 mg/ml, a sustained elevation in  $[Ca^{2+}]_c$  was observed (Figure 2A). Our data demonstrated that  $[Ca^{2+}]_c$  increased as a function of  $TiO_2$  concentration and coincided with the measurements of histamine secretion (Figure 5A). The  $TiO_2$  NPs used in our study consisted of a mixture of anatase/rutile crystalline forms which have been shown to impose greater cellular toxicity [22]. Although phase composition, surface chemistry and purity may play important roles in influencing nanotoxicity, dependence on additional physical parameters such as particle size, surface area and concentration have all been reported [20,21,36-38]. However, the mechanism by which  $TiO_2$  NPs trigger intracellular mast cell  $Ca^{2+}$  signaling is not clear.

Previous experimental evidence suggests that ultrafine carbon black and ZnO NPs can induce extracellular  $Ca^{2+}$  influx through activated voltage-gated  $Ca^{2+}$  channels [35,39]. The question of whether  $TiO_2$  NPs can trigger a similar process is unclear. Our experiments confirmed that  $[Ca^{2+}]_c$  failed to increase significantly when cells were treated with  $TiO_2$  NPs in  $Ca^{2+}$ -free Hanks' solution (Figure 2B) or blocked with  $CdCl_2$  (a general  $Ca^{2+}$

channel blocker) (Figure 2C) [17]. We then demonstrated that  $TiO_2$  NPs can activate L-type voltage-gated  $Ca^{2+}$  channels on mast cells, allowing extracellular  $Ca^{2+}$  influx into the cytosol (Figure 2D and 2E). The ROS generated in response to NP exposure has been attributed to the opening of membrane  $Ca^{2+}$  channels [17,35,39,40]. Our data revealed that  $TiO_2$  NPs elevated cytosolic ROS levels (Figure 3A) and that the involvement of oxidative stress could modulate intracellular  $Ca^{2+}$  homeostasis by activating L-type  $Ca^{2+}$  channels (Figure 3B). Supporting our results, Hussain and colleagues also reported that  $TiO_2$  NPs can exert oxidative effects under abiotic conditions [23]. Other studies have shown that  $TiO_2$  NPs increase intracellular ROS production in various cells by causing mitochondrial injuries, such as impaired mitochondrial membrane permeability, a decrease in mitochondrial potential, mitochondrial respiratory dysfunction and downregulation of SOD (superoxide dismutase) and GSH (glutathione) levels [41,42]. ROS are thought to affect  $Ca^{2+}$  signaling possibly by oxidation of thiol groups of membrane channels and changing the intra-molecular bonding of proteins and lipids [43,44]. Moreover, it has been suggested that ROS can impact membrane  $Ca^{2+}$  channels and  $Ca^{2+}$  binding proteins [43]. In addition to ROS formation, direct damage to cell membrane integrity by lipid peroxidation may be an alternative route for extracellular  $Ca^{2+}$  entry [22,39]. Co-administration of  $TiO_2$  NPs and fluorescent calcein dye resulted in an augmented calcein permeation ratio (Figure 2F). The calcein data corroborates with previous publications showing that  $TiO_2$  NPs can perturb the lipid bilayer, perhaps by forming transient pores that may account for the portion of  $Ca^{2+}$  increase that could not be completely abolished by blocking L-type  $Ca^{2+}$  channels [17,45].




**Figure 4** Measurement of  $[Ca^{2+}]_c$  after stimulation by  $TiO_2$  NPs. RBL-2H3 cells were stimulated with  $TiO_2$  NPs with concentrations that vary from 0.1 mg/ml-1 mg/ml, in the presence of (A) U73122 (10  $\mu$ M), (B) 2-APB (50  $\mu$ M) and (C) Xestospongine (20  $\mu$ M) and (D) Thapsigargin (100 nM). The colors used are in accordance with Figure 2A. Each line represents the average fluorescent intensity of more than 200 cells.

An initial upsurge in  $[Ca^{2+}]_c$  from extracellular influx is usually relayed by a secondary  $Ca^{2+}$  release from internal organelles [46]. However, the question of how  $TiO_2$  NPs transduce  $Ca^{2+}$  signals in mast cells involving common secondary messengers remains unanswered. Heretofore, we demonstrated that  $TiO_2$  NPs could stimulate PLC and  $IP_3$  receptor activities in mast cells, resulting in secondary amplification of cytosolic  $Ca^{2+}$ . Data from Figure 4A-C showed that cells incubated with U73122, 2-APB and Xestospongine notably inhibited  $TiO_2$  NP-triggered  $Ca^{2+}$  rise and histamine secretion (Figure 5B). Activation of PLC can be explained by the stimulation from ROS [47,48]. Alternatively, a localized increase in  $[Ca^{2+}]_c$  has been shown to trigger PLC, thereby generating  $IP_3$  and DAG [49]. Collectively, these results point toward the contribution from ER

$Ca^{2+}$  stores. Figure 4D revealed that depleting the ER internal store with Thapsigargin significantly diminished  $TiO_2$  NPs stimulated cytosolic  $Ca^{2+}$  increase and subsequent histamine secretion (Figure 5B). Our data indicates that the ER dependent  $IP_3$ - $IP_3$  receptor pathway is critically involved in amplifying (or sustaining) the initial cytosolic  $Ca^{2+}$  rise and subsequent histamine release.

### Conclusion

Our findings revealed a new mechanism depicting how  $TiO_2$  NPs can potentially exacerbate many allergic inflammatory responses and perhaps non-allergic inflammatory disorders. Mast cell exposure to  $TiO_2$  NPs can activate membrane L-type  $Ca^{2+}$  channels, induce ROS production and stimulate PLC activity. Influx of extracellular  $Ca^{2+}$



**Figure 5 Measurement of histamine secretion evoked by TiO<sub>2</sub> NPs.** (A) ELISA quantification of histamine release from RBL-2H3 cells after TiO<sub>2</sub> NP (0.1 mg/ml-0.75 mg/ml stimulation) in normal Hanks' solution (n ≥ 3, \*\*p < 0.005). (B) Subsequent assessment of histamine secretion from RBL-2H3 cells when pre-treated with an intracellular Ca<sup>2+</sup> chelator (BAPTA) and various blockers prior to TiO<sub>2</sub> NP exposure (n ≥ 3; white, checkboard, and black bars represent 0, 0.5 and 0.75 mg/ml, respectively).

raises [Ca<sup>2+</sup>]<sub>C</sub>, and when coupled with the IP<sub>3</sub>-IP<sub>3</sub> receptor pathway, can trigger the release of ER resident Ca<sup>2+</sup> and subsequent histamine secretion. These results suggest that mast cell degranulation of histamine may be significantly augmented and intensified in NP exposed tissues with or without IgE antibody-based sensitization. This model may also provide a new explanation for chronic inflammatory diseases elicited by airborne PM. The inhaled NPs can circulate and accumulate in various organs thereby elevating the risks of activating mast cell degranulation in other tissues [25,50,51]. As a result, NP exposure may worsen the mast cell associated inflammatory symptoms involving arthritis, atherosclerosis and coronary diseases [52-54]. Finally, our results suggest that a new immunoregulatory system may be considered since NPs can directly trigger inflammatory mediators, thereby bypassing traditional immuno-stimulation by allergens.

## Methods

### Culture of RBL-2H3 Mast Cells

The RBL-2H3 rat mast cell line (ATCC, Manassas, VA, USA) is a widely-used mast cell model that responds to stimuli by secreting histamine and other mediators [55]. RBL-2H3 cells elicit a potent immune allergic response following crosslinking of their IgE-bound FcεRI by multivalent allergens [55]. Cells were cultured in 15 cm cell culture plates (VWR, CA, USA) in MEM medium (Invitrogen, CA, USA) supplemented with L-glutamine, 1% penicillin/streptomycin and 10% heat inactivated fetal bovine serum (FBS) (Invitrogen, CA, USA). Cultures were incubated in a humidified incubator at 37°C/5% CO<sub>2</sub>. Cell counts were performed using trypan blue

(Sigma-Aldrich, MO, USA) exclusion and a Bright-Line haemocytometer.

### Nanoparticles and Characterization

A mixture of anatase and rutile forms of ultrafine titanium (IV) dioxide (< 100 nm diameter by Brunauer Emmett Teller (BET) method, purity of 99.5% trace metals basis) (Cat No. 634662) (Sigma-Aldrich, MO, USA) was used in this study because this form has been shown to result in more severe cellular injuries [22]. The TiO<sub>2</sub> NPs have a primary crystalline size of 100 nm (maximum), specific surface area of 46.3 m<sup>2</sup>/g (as determined by BET), equivalent spherical diameter of 100 nm (maximum) and a trace metallic impurities of 1000 ppm (maximum) (information from Sigma). All TiO<sub>2</sub> NP samples were reconstituted in Hanks' buffer (Invitrogen, CA, USA) and sonicated for approximately 1 minute immediately before usage. The concentrations used were 1 mg/ml, 0.75 mg/ml, 0.5 mg/ml, 0.25 mg/ml, and 0.1 mg/ml. All sizes of NP suspension were independently confirmed using homodyne dynamics laser scattering (DLS) as described in the previous study [17]. Morphology and size of TiO<sub>2</sub> NP powder were determined by transmission electron microscope (JEOL JEM-2010 HRTEM, MA, USA) and analyzed using an analysisPRO (Olympus, PA, USA).

### Cell Preparation

Cells were seeded at 1 × 10<sup>5</sup> cells per well in a 24-well plate, and incubated for 24 hrs in MEM medium supplemented with 10% FBS. Following 24 hr incubation, the MEM medium was removed from the cells and the

culture was rinsed with Hanks' solution twice before use.

#### Measurement of cytosolic $\text{Ca}^{2+}$ concentrations induced by $\text{TiO}_2$ exposure

All experiments were performed in dark conditions. The cells were loaded with a Rhod-2 AM dye (1  $\mu\text{M}$ ) ( $K_d = 570$  nM,  $\lambda_{\text{Ex}} = 552$  nm and  $\lambda_{\text{Em}} = 581$ ) (Invitrogen, CA, USA) for 45 minutes. After the dye loading, the cells were rinsed and incubated with either normal Hanks' or  $\text{Ca}^{2+}$ -free Hanks' solution, and treated with the appropriate  $\text{TiO}_2$  NP concentrations. All  $\text{Ca}^{2+}$  signaling experiments were carried out on a thermoregulated stage at  $37^\circ\text{C}$  mounted on a Nikon microscope (Nikon Eclipse TE2000-U, Tokyo, Japan). RBL-2H3 cells were then incubated with cadmium chloride (200  $\mu\text{M}$ ; Sigma-Aldrich, MO, USA) to block the membrane  $\text{Ca}^{2+}$  channels [17], followed by  $\text{TiO}_2$  NP stimulation. To test the interaction between  $\text{TiO}_2$  and L-type membrane  $\text{Ca}^{2+}$  channels, nifedipine (10  $\mu\text{M}$ ; Sigma-Aldrich, MO, USA) and verapamil (100  $\mu\text{M}$ ; Sigma-Aldrich, MO, USA), L-type  $\text{Ca}^{2+}$  channel blockers [17,39] were applied to RBL-2H3 cells, independently, prior to the exposure of  $\text{TiO}_2$ . Antioxidant N-acetylcysteine (NAC, 250  $\mu\text{M}$ ; Sigma-Aldrich, MO, USA) was also added to RBL-2H3 cells to study the involvement of reactive oxygen species (ROS) [17,39], possibly generated as a result of  $\text{TiO}_2$  stimulation, and the activation of  $\text{Ca}^{2+}$  channels. Thapsigargin (100 nM; Sigma-Aldrich, MO, USA) was used to deplete the ER  $\text{Ca}^{2+}$  content in order to investigate the contribution of the internal ER  $\text{Ca}^{2+}$  pool [17,56]. In order to test the involvement of secondary messenger molecules in intensifying a  $\text{Ca}^{2+}$  response, U73122 (PLC blocker) (10  $\mu\text{M}$ ; Sigma-Aldrich, MO, USA), Aminoethoxydiphenyl borate (2-APB) (50  $\mu\text{M}$ ; Sigma-Aldrich, MO, USA) and Xestospongins C (IP<sub>3</sub> receptor blockers) (20  $\mu\text{M}$ ; WVR, CA, USA) were pre-incubated with cells prior to  $\text{TiO}_2$  NP exposure [56,57].

#### Calcein dye leakage measurements

RBL-2H3 cells were seeded at a density of  $1 \times 10^5$  cells per well in a 24-well plate and cultured for 24 hrs.  $\text{TiO}_2$  prepared with calcein fluorescent dye (50  $\mu\text{M}$ ) (Invitrogen, CA, USA) in Hanks' buffer was incubated with the cells for 5 minutes at  $37^\circ\text{C}$ . Calcein is a biological inert green-fluorescent molecule of a molecular mass of 623 Daltons and an estimated molecular radius of 0.6 nm [17]. The  $\text{TiO}_2$  NP solution containing calcein dye was then removed and the cells were rinsed twice with PBS to remove any possible remnants of the dye. Subsequently, the cells were loaded with Hoechst (10  $\mu\text{M}$ ) (Sigma-Aldrich, MO, USA), a fluorescent nucleus dye, for 5 minutes at  $37^\circ\text{C}$ , and were then rinsed thoroughly [17]. Fresh Hanks' solution was added into each well before taking

fluorescent images of calcein and Hoechst loaded cells with a Nikon fluorescence microscope. Afterwards, a percentage of calcein loaded cells against total number of cells, as determined by the Hoechst dye, was calculated for each of the  $\text{TiO}_2$  NP concentrations used in the experiment.

#### Intracellular reactive oxygen species (ROS) production

ROS production was evaluated by fluorescence microscopy using oxidation of CM-H<sub>2</sub>DCFDA dye (Invitrogen, CA, USA). The cells ( $1 \times 10^5$  cells/well) were cultured for 24 hrs before being rinsed with PBS solution. Hanks' buffer, containing  $\text{TiO}_2$  NPs at concentrations ranging from 0-0.75 mg/ml, was then incubated with the cells for 15 minutes in  $37^\circ\text{C}$  followed by PBS washing and loading with 2  $\mu\text{M}$  CM-H<sub>2</sub>DCFDA dye for 30 minutes. Fluorescent images of ROS generated in cells were captured and analyzed by calculating the ratio of increase in fluorescent intensity between  $\text{TiO}_2$  NP treatment and control groups.

#### Histamine detection with enzyme linked immunosorbent assay (ELISA) preparation

The cells were seeded at a density of  $1 \times 10^5$  cells density in a 24-well plate and cultured for 24 hrs. RBL-2H3 cells were then rinsed with PBS and pre-treated with BAPTA-AM (50  $\mu\text{M}$ ) (Invitrogen, CA, USA), Thapsigargin (SERCA pump inhibitor), U73122, 2-APB, or Xestospongins for 20 minutes in the same manner. Cells were stimulated for 5 minutes with the appropriate  $\text{TiO}_2$  NP concentrations (0-0.75 mg/ml) which were prepared in PBS. The histamine-containing supernatant was collected and centrifuged to remove any remaining  $\text{TiO}_2$  NPs. The supernatant was then incubated in a 96-well plate overnight at  $4^\circ\text{C}$ . The rest of the assay was carried out in accord to Neogen ELISA instructions pertaining to the histamine kit (Neogen Corp, MI, USA).

#### Image Analysis

After staining the treated cells, image analysis was performed with an inverted Nikon Eclipse TE2000-U fluorescent microscope. Each photo was taken at a magnification of  $200 \times$  and analyzed using SimplePCI (Compix Inc., Imaging Systems, Sewickley, PA, USA). The data shown is a representative of  $\text{Ca}^{2+}$  signals of more than 200 cells and the experiments were conducted independently for at least 3 times.

#### Statistical Analysis

The data was presented as means  $\pm$  SD. Each experiment was performed independently at least three times. Statistical significance was determined using a Student's t-test analysis with p values  $< 0.05$  (GraphPad Prism 4.0, GraphPad Software, Inc., San Diego, CA, USA).



### Acknowledgements

The authors gratefully thank David Ojcius for careful reading of the manuscript. We also thank Mike Dunlap from IMF for assistance with TEM imaging. This study was supported by grants from NIH (1R15HL095039), NSF (CBET-0932404) and the UC CITRIS Program. EYC, MG and CSC were supported by UC Merced GRC summer fellowships, NIH NHLBI URS and Center of Excellence on Health Disparities (1P20MD005049-01 from the National Center on Minority Health and Health Disparities).

### Authors' contributions

EYC, MG, YCW: Designed research, conducted experiments, analyzed and interpreted data. EYC, MG, YCW, AJM, CSC and WCC: wrote, reviewed and revised the manuscript. All authors have read and approved the final manuscript.

### Competing interests

The authors declare that they have no competing interests.

Received: 22 July 2011 Accepted: 19 January 2012

Published: 19 January 2012

### References

- Galli SJ, Tsai M, Piliponsky AM: The development of allergic inflammation. *Nature* 2008, **454**:445-454.
- Jones BL, Kearns GL: Histamine: new thoughts about a familiar mediator. *Clin Pharmacol Ther* 2011, **89**:189-197.
- Williams JL, Gow JA, Klier SM, McCue SL, Salapatek AM, McNamara TR: Non-clinical pharmacology, pharmacokinetics, and safety findings for the antihistamine bepotastine besilate. *Curr Med Res Opin* 2010, **26**:2329-2338.
- Jutel M, Akdis M, Akdis CA: Histamine, histamine receptors and their role in immune pathology. *Clin Exp Allergy* 2009, **39**:1786-1800.
- Holgate ST: The epidemic of allergy and asthma. *Nature* 1999, **402**:B2-4.
- von Klot S, Wolke G, Tuch T, Heinrich J, Dockery DW, Schwartz J, Kreyling WG, Wichmann HE, Peters A: Increased asthma medication use in association with ambient fine and ultrafine particles. *Eur Respir J* 2002, **20**:691-702.
- Spira-Cohen A, Chen LC, Kendall M, Lall R, Thurston GD: Personal Exposures to Traffic-Related Air Pollution and Acute Respiratory Health Among Bronx School Children with Asthma. *Environ Health Perspect* 2011.
- Downs SH, Schindler C, Liu LJ, Keidel D, Bayer-Oglesby L, Brutsche MH, Gerbase MW, Keller R, Kunzli N, Leuenberger P, et al: Reduced exposure to PM10 and attenuated age-related decline in lung function. *N Engl J Med* 2007, **357**:2338-2347.
- Archer AJ, Cramton JL, Pfau JC, Colasurdo G, Holian A: Airway responsiveness after acute exposure to urban particulate matter 1648 in a DO11.10 murine model. *Am J Physiol Lung Cell Mol Physiol* 2004, **286**:L337-343.
- Hussain S, Vanoirbeek JA, Luyts K, De Vooght V, Verbeken E, Thomassen LC, Martens JA, Dinsdale D, Boland S, Marano F, et al: Lung exposure to nanoparticles modulates an asthmatic response in a mouse model of asthma. *Eur Respir J* 2011.
- Jin C, Shelburne CP, Li G, Potts EN, Riebe KJ, Sempowski GD, Foster WM, Abraham SN: Particulate allergens potentiate allergic asthma in mice through sustained IgE-mediated mast cell activation. *J Clin Invest* 2011.
- Yanagisawa R, Takano H, Inoue K, Koike E, Kamachi T, Sadakane K, Ichinose T: Titanium dioxide nanoparticles aggravate atopic dermatitis-like skin lesions in NC/Nga mice. *Exp Biol Med (Maywood)* 2009, **234**:314-322.
- Hugg TT, Jaakkola MS, Ruotsalainen RO, Pushkarev VJ, Jaakkola JJ: Parental smoking behaviour and effects of tobacco smoke on children's health in Finland and Russia. *Eur J Public Health* 2008, **18**:55-62.
- Simoni M, Baldacci S, Puntoni R, Pistelli F, Farchi S, Lo Presti E, Pistelli R, Corbo G, Agabiti N, Basso S, et al: Respiratory symptoms/diseases and environmental tobacco smoke (ETS) in never smoker Italian women. *Respir Med* 2007, **101**:531-538.
- Kulig M, Luck W, Lau S, Niggemann B, Bergmann R, Klettke U, Guggenmoos-Holzmann I, Wahn U: Effect of pre- and postnatal tobacco smoke exposure on specific sensitization to food and inhalant allergens during the first 3 years of life. Multicenter Allergy Study Group, Germany. *Allergy* 1999, **54**:220-228.
- Kaida T, Kobayashi K, Adachi M, Suzuki F: Optical characteristics of titanium oxide interference film and the film laminated with oxides and their applications for cosmetics. *J Cosmet Sci* 2004, **55**:219-220.
- Chen EY, Garnica M, Wang YC, Chen CS, Chin WC: Mucin secretion induced by titanium dioxide nanoparticles. *PLoS One* 2011, **6**:e16198.
- Hussain S, Thomassen LC, Ferecatu I, Borot MC, Andreau K, Martens JA, Fleury J, Baeza-Squiban A, Marano F, Boland S: Carbon black and titanium dioxide nanoparticles elicit distinct apoptotic pathways in bronchial epithelial cells. *Part Fibre Toxicol* 2010, **7**:10.
- Ahn MH, Kang CM, Park CS, Park SJ, Rhim T, Yoon PO, Chang HS, Kim SH, Kyono H, Kim KC: Titanium dioxide particle-induced goblet cell hyperplasia: association with mast cells and IL-13. *Respir Res* 2005, **6**:34.
- Sayes CM, Wahi R, Kurian PA, Liu Y, West JL, Ausman KD, Warheit DB, Colvin VL: Correlating nanoscale titania structure with toxicity: a cytotoxicity and inflammatory response study with human dermal fibroblasts and human lung epithelial cells. *Toxicol Sci* 2006, **92**:174-185.
- Jiang J, Oberdorster G, Elder A, Gelein R, Mercer P, Biswas P: Does Nanoparticle Activity Depend upon Size and Crystal Phase? *Nanotoxicology* 2008, **2**:33-42.
- Gurr JR, Wang AS, Chen CH, Jan KY: Ultrafine titanium dioxide particles in the absence of photoactivation can induce oxidative damage to human bronchial epithelial cells. *Toxicology* 2005, **213**:66-73.
- Hussain S, Boland S, Baeza-Squiban A, Hamel R, Thomassen LC, Martens JA, Billon-Galland MA, Fleury-Feith J, Moisan F, Pairon JC, Marano F: Oxidative stress and proinflammatory effects of carbon black and titanium dioxide nanoparticles: role of particle surface area and internalized amount. *Toxicology* 2009, **260**:142-149.
- Pujalte I, Passagne I, Brouillaud B, Treguer M, Durand E, Ohayon-Courtes C, L'Azou B: Cytotoxicity and oxidative stress induced by different metallic nanoparticles on human kidney cells. *Part Fibre Toxicol* 2011, **8**:10.
- Choi HS, Ashitate Y, Lee JH, Kim SH, Matsui A, Insin N, Bawendi MG, Semmler-Behnke M, Frangioni JV, Tsuda A: Rapid translocation of nanoparticles from the lung airspaces to the body. *Nat Biotechnol* 2010, **28**:1300-1303.
- Li Y, Li J, Yin J, Li W, Kang C, Huang Q, Li Q: Systematic influence induced by 3 nm titanium dioxide following intratracheal instillation of mice. *J Nanosci Nanotechnol* 2010, **10**:8544-8549.
- Oberdorster G, Sharp Z, Atudorei V, Elder A, Gelein R, Lunts A, Kreyling W, Cox C: Extrapulmonary translocation of ultrafine carbon particles following whole-body inhalation exposure of rats. *J Toxicol Environ Health A* 2002, **65**:1531-1543.
- Barlow PG, Clouter-Baker A, Donaldson K, MacCallum J, Stone V: Carbon black nanoparticles induce type II epithelial cells to release chemotaxins for alveolar macrophages. *Part Fibre Toxicol* 2005, **2**:11.
- Chen HW, Su SF, Chien CT, Lin WH, Yu SL, Chou CC, Chen JJ, Yang PC: Titanium dioxide nanoparticles induce emphysema-like lung injury in mice. *FASEB J* 2006, **20**:2393-2395.
- Moon C, Park HJ, Choi YH, Park EM, Castranova V, Kang JL: Pulmonary inflammation after intraperitoneal administration of ultrafine titanium dioxide (TiO<sub>2</sub>) at rest or in lungs primed with lipopolysaccharide. *J Toxicol Environ Health A* 2010, **73**:396-409.
- Oberdorster G, Ferin J, Gelein R, Soderholm SC, Finkelstein J: Role of the alveolar macrophage in lung injury: studies with ultrafine particles. *Environ Health Perspect* 1992, **97**:193-199.
- Oberdorster G, Ferin J, Lehnert BE: Correlation between particle size, in vivo particle persistence, and lung injury. *Environ Health Perspect* 1994, **102**(Suppl 5):173-179.
- Abdullah LH, Conway JD, Cohn JA, Davis CW: Protein kinase C and Ca<sup>2+</sup> activation of mucin secretion in airway goblet cells. *Am J Physiol* 1997, **273**:L201-210.
- Petersen CC, Toescu EC, Petersen OH: Different patterns of receptor-activated cytoplasmic Ca<sup>2+</sup> oscillations in single pancreatic acinar cells: dependence on receptor type, agonist concentration and intracellular Ca<sup>2+</sup> buffering. *Embo J* 1991, **10**:527-533.
- Stone V, Tuinman M, Vamvakopoulos JE, Shaw J, Brown D, Petterson S, Faux SP, Borm P, MacNee W, Michaelangeli F, Donaldson K: Increased calcium influx in a monocytic cell line on exposure to ultrafine carbon black. *Eur Respir J* 2000, **15**:297-303.
- Kobayashi N, Naya M, Endoh S, Maru J, Yamamoto K, Nakanishi J: Comparative pulmonary toxicity study of nano-TiO<sub>2</sub> particles of

- different sizes and agglomerations in rats: different short- and long-term post-instillation results. *Toxicology* 2009, **264**:110-118.
37. Oberdorster G, Oberdorster E, Oberdorster J: **Nanotoxicology: an emerging discipline evolving from studies of ultrafine particles.** *Environ Health Perspect* 2005, **113**:823-839.
  38. Monteiller C, Tran L, MacNee W, Faux S, Jones A, Miller B, Donaldson K: **The pro-inflammatory effects of low-toxicity low-solubility particles, nanoparticles and fine particles, on epithelial cells in vitro: the role of surface area.** *Occup Environ Med* 2007, **64**:609-615.
  39. Huang CC, Aronstam RS, Chen DR, Huang YW: **Oxidative stress, calcium homeostasis, and altered gene expression in human lung epithelial cells exposed to ZnO nanoparticles.** *Toxicol In Vitro* 2009, **24**:45-55.
  40. Donaldson K, Stone V, Borm PJ, Jimenez LA, Gilmour PS, Schins RP, Knaapen AM, Rahman I, Faux SP, Brown DM, MacNee W: **Oxidative stress and calcium signaling in the adverse effects of environmental particles (PM10).** *Free Radic Biol Med* 2003, **34**:1369-1382.
  41. Shukla RK, Sharma V, Pandey AK, Singh S, Sultana S, Dhawan A: **ROS-mediated genotoxicity induced by titanium dioxide nanoparticles in human epidermal cells.** *Toxicol In Vitro* 2011, **25**:231-241.
  42. Freyre-Fonseca V, Delgado-Buenrostro NL, Gutierrez-Cirlos EB, Calderon-Torres CM, Cabellos-Avelar T, Sanchez-Perez Y, Pinzon E, Torres I, Molina-Jijon E, Zazueta C, et al: **Titanium dioxide nanoparticles impair lung mitochondrial function.** *Toxicol Lett* 2011, **202**:111-119.
  43. Kourie JI: **Interaction of reactive oxygen species with ion transport mechanisms.** *Am J Physiol* 1998, **275**:C1-24.
  44. Palaniappan PR, Pramod KS: **The effect of titanium dioxide on the biochemical constituents of the brain of Zebrafish (Danio rerio): An FT-IR study.** *Spectrochim Acta A Mol Biomol Spectrosc* 2011.
  45. Kelly CV, Leroueil PR, Orr BG, Banaszak Holl MM, Andricioaei I: **Poly (amidoamine) dendrimers on lipid bilayers II: Effects of bilayer phase and dendrimer termination.** *J Phys Chem B* 2008, **112**:9346-9353.
  46. Berridge MJ, Irvine RF: **Inositol trisphosphate, a novel second messenger in cellular signal transduction.** *Nature* 1984, **312**:315-321.
  47. Lu SP, Lin Feng MH, Huang HL, Huang YC, Tsou WI, Lai MZ: **Reactive oxygen species promote raft formation in T lymphocytes.** *Free Radic Biol Med* 2007, **42**:936-944.
  48. Muller G, Wied S, Jung C, Over S: **Hydrogen peroxide-induced translocation of glycolipid-anchored (c)AMP-hydrolases to lipid droplets mediates inhibition of lipolysis in rat adipocytes.** *Br J Pharmacol* 2008, **154**:901-913.
  49. Ashby MC, Craske M, Park MK, Gerasimenko OV, Burgoyne RD, Petersen OH, Tepikin AV: **Localized Ca<sup>2+</sup> uncaging reveals polarized distribution of Ca<sup>2+</sup>-sensitive Ca<sup>2+</sup> release sites: mechanism of unidirectional Ca<sup>2+</sup> waves.** *J Cell Biol* 2002, **158**:283-292.
  50. Kumar R, Roy I, Ohulchanskyy TY, Vathy LA, Bergey EJ, Sajjad M, Prasad PN: **In vivo biodistribution and clearance studies using multimodal organically modified silica nanoparticles.** *ACS Nano* 2010, **4**:699-708.
  51. Semmler-Behnke M, Kreyling WG, Lipka J, Fertsch S, Wenk A, Takenaka S, Schmid G, Brandau W: **Biodistribution of 1.4- and 18-nm gold particles in rats.** *Small* 2008, **4**:2108-2111.
  52. Woolley DE: **The mast cell in inflammatory arthritis.** *N Engl J Med* 2003, **348**:1709-1711.
  53. Hansson GK, Robertson AK, Soderberg-Naucler C: **Inflammation and atherosclerosis.** *Annu Rev Pathol* 2006, **1**:297-329.
  54. Ozben B, Erdogan O: **The role of inflammation and allergy in acute coronary syndromes.** *Inflamm Allergy Drug Targets* 2008, **7**:136-144.
  55. Passante E, Frankish N: **The RBL-2H3 cell line: its provenance and suitability as a model for the mast cell.** *Inflamm Res* 2009, **58**:737-745.
  56. Nguyen T, Chin WC, Verdugo P: **Role of Ca<sup>2+</sup>/K<sup>+</sup> ion exchange in intracellular storage and release of Ca<sup>2+</sup>.** *Nature* 1998, **395**:908-912.
  57. Divangahi M, Balghi H, Danialou G, Comtois AS, Demoule A, Ernest S, Haston C, Robert R, Hanrahan JW, Radzioch D, Petrof BJ: **Lack of CFTR in skeletal muscle predisposes to muscle wasting and diaphragm muscle pump failure in cystic fibrosis mice.** *PLoS Genet* 2009, **5**:e1000586.

doi:10.1186/1743-8977-9-2

**Cite this article as:** Chen et al.: A mixture of anatase and rutile TiO<sub>2</sub> nanoparticles induces histamine secretion in mast cells. *Particle and Fibre Toxicology* 2012 **9**:2.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

